

# Luminescence dating of young volcanic eruptions

Tuesday, 11 February 2020 16:00 (2 minutes)

Reliable chronologies of volcanic eruptions are vital for hazard analysis, but dating of Holocene and Late Pleistocene volcanism poses a major challenge. Established techniques such as  $^{40}\text{Ar}/^{39}\text{Ar}$  are often problematic due to the long half-life of  $^{40}\text{K}$  or the absence of datable materials. In this context, luminescence dating is a promising alternative.

Luminescence signal resetting during volcanic activity can be caused by heat (lava, contact to lava), light (disintegration of ejecta) or (temperature-assisted) pressure in the course of phreatomagmatic explosions. While volcanogenic minerals forming volcanic rocks are less suitable for luminescence dating due to anomalous fading, the signal of volcanogenically heated or fragmented country rock also relates to the time of eruption and further provides more accurate results, due to the presence of minerals like quartz.

This contribution aims to illustrate the potential of this latter approach by presenting two case studies. The first refers to two Late Pleistocene scoria cones in the Westeifel Volcanic Field (WEVF), Germany, of which the Wartgesberg locality was dated by  $^{40}\text{Ar}/^{39}\text{Ar}$  and  $^{14}\text{C}$ , while the closeby Facher Höhe is chronologically poorly constrained [1]. The former locality allows testing the accuracy of various luminescence techniques (TL, OSL, IRSL) applied to quartz and polymineral separates against independent age control. The other study site is the monogenetic Lake Nyos Maar, which is part of the Cameroon Volcanic Line (CVL). Previous dating efforts of the last explosive activity are inconsistent and yielded age estimates ranging from 400 a ( $^{14}\text{C}$ ) to >350 ka (K-Ar) [2].

Our results demonstrate that multiple luminescence methods (TL, OSL) yield equally valid age estimates averaging to  $33.6 \pm 2.4$  ka for the Wartgesberg site, which is in good agreement with  $^{40}\text{Ar}/^{39}\text{Ar}$  and  $^{14}\text{C}$  results. The Facher Höhe, however, is much younger than previously assumed with an average TL age of  $15.5 \pm 1.1$  ka. This southeastern part of the WEVF thus hosts many of the most recent eruption sites, which has important implications for modelling the causes and timing of Eifel volcanism and assessment of future eruption locations. Luminescence studies for the Nyos Maar suggest that the phreatomagmatic explosion has completely reset the inherited TL signal and an average maximum age of  $12.3 \pm 1.5$  ka roughly fits with U-series age estimates [2], indicating that TL is a viable tool for completing the fragmentary chronostratigraphic record of the CVL.

## References

1. Mertz, D.F., Löhnertz, W., Nomade, S., Pereira, A., Prelevic, D., Renne, P.R., 2015. Temporal-spatial evolution of low-SiO<sub>2</sub> volcanism in the Pleistocene West Eifel volcanic field (West Germany) and relationship to upwelling asthenosphere. *J. Geodyn.* 88, 59-79.
2. Aka, F.T., Yokoyama, T., Kusakabe, M., Nakamura, E., Tanyileke, G., Ateba, B., Ngako, V., Nnange, J., Hell, J., 2008. U-series dating of Lake Nyos maar basalts, Cameroon (West Africa): Implications for potential hazards on the Lake Nyos dam. *J. Volcanol. Geoth. Res.* 176, 212-224.

**Primary author:** SCHMIDT, Christoph

**Co-authors:** Mrs SCHAARSCHMIDT, Maria (University of Wollongong); Dr KOLB, Thomas (JLU Gießen); Dr RICHTER, Daniel (Max Planck Institute for Evolutionary Anthropology, Leipzig); Prof. TCHOUANKOUE, Jean Pierre (University of Yaounde I); Prof. ZÖLLER, Ludwig (University of Bayreuth)

**Presenter:** SCHMIDT, Christoph

**Session Classification:** Monitoring and Risk Assessment

**Track Classification:** Monitoring and Risk Assessment