

## Online adaptive estimation of decoherence timescales for a single qubit

*Monday 19 September 2022 12:15 (15 minutes)*

The estimation of decoherence timescales is important not only as a key performance indicator for quantum technology, but also to measure physical quantities through the change they induce in the relaxation of quantum sensors. Typically, decoherence times are estimated by fitting a signal acquired while sweeping the time delay between qubit preparation and detection on a pre-determined range. Here we describe an adaptive Bayesian approach, based on a simple analytical update rule, to estimate  $T_1$ ,  $T_2^*$  and  $T_2$  with the fewest number of measurements, demonstrating a speed-up of factor 3-10, depending on the specific experiment, compared to the standard protocols. We also demonstrate that, when sensing time  $\tau$  is the resource to be minimised, a further speed-up of a factor  $\sim 2$  can be obtained by maximising the ratio between Fisher information and time  $\tau$ , compared to the Fisher information.

We demonstrate the online adaptive protocols on a single electronic spin qubit associated with a nitrogen-vacancy (NV) centre in diamond, implementing Bayesian inference on a hard-realtime microcontroller in less than 100  $\mu$ s, a time negligible compared to the duration of each measurement. Our protocol can be readily applied to different types of quantum systems.

**Author:** ARSHAD, Muhammad Junaid (Heriot-Watt University)

**Co-authors:** Mr BEKKER, Christiaan (Heriot Watt University); Mr HAYLOCK, Ben (Heriot Watt University); Mr SKRZYPCZAK, Krzysztof (Heriot Watt University); Mr WHITE, Daniel (Heriot Watt University); Mr GAUGER, Erik (Heriot Watt University); Mr BONATO, Cristian (Heriot Watt University)

**Presenter:** ARSHAD, Muhammad Junaid (Heriot-Watt University)

**Session Classification:** Optimization and Control

**Track Classification:** Optimization and Control